



**QUEEN'S
UNIVERSITY
BELFAST**

Assessing elements of a family approach to reduce adolescent drinking frequency: parent-adolescent relationship, knowledge management, and keeping secrets

McCann, M., Perra, O., McLaughlin, A., McCartan, C., & Higgins, K. (2016). Assessing elements of a family approach to reduce adolescent drinking frequency: parent-adolescent relationship, knowledge management, and keeping secrets. *Addiction*, 111(5), 843-853. <https://doi.org/10.1111/add.13258>

Published in:
Addiction

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2015 The Authors.

This is an open access article under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Assessing elements of a family approach to reduce adolescent drinking frequency: parent–adolescent relationship, knowledge management and keeping secrets

Mark McCann¹, Oliver Perra², Aisling McLaughlin³, Claire McCartan³ & Kathryn Higgins³

MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, UK,¹ School of Nursing and Midwifery, Queen's University Belfast, Belfast, UK² and Institute of Child Care Research, Queen's University Belfast, Belfast, UK³

ABSTRACT

Aims To estimate (1) the associations between parent–adolescent relationship, parental knowledge and subsequent adolescent drinking frequency and (2) the influence of alcohol use on parental knowledge. **Design** Path analysis of school based cohort study with annual surveys. **Setting** Post-primary schools from urban and intermediate/rural areas in Northern Ireland. **Participants** A total of 4937 post-primary school students aged approximately 11 years in 2000 followed until approximately age 16 years in 2005. **Measurements** Pupil-reported measures of: frequency of alcohol use; parent–child relationship quality; subdimensions of parental monitoring: parental control, parental solicitation, child disclosure and child secrecy. **Findings** Higher levels of parental control [ordinal logistic odds ratio (OR) = 0.86, 95% confidence interval (CI) = 0.78, 0.95] and lower levels of child secrecy (OR = 0.83, 95% CI = 0.75, 0.92) were associated subsequently with less frequent alcohol use. Parental solicitation and parent–child relationship quality were not associated with drinking frequency. Weekly alcohol drinking was associated with higher subsequent secrecy (beta = −0.42, 95% CI = −0.53, −0.32) and lower parental control (beta = −0.15, 95% CI = −0.26, −0.04). Secrecy was more strongly predictive of alcohol use at younger compared with older ages ($P = 0.02$), and alcohol use was associated less strongly with parental control among families with poorer relationships ($P = 0.04$). **Conclusions** Adolescent alcohol use appears to increase as parental control decreases and child secrecy increases. Greater parental control is associated with less frequent adolescent drinking subsequently, while parent–child attachment and parental solicitation have little influence on alcohol use.

Keywords Adolescence, alcohol, child, parental monitoring, parent–child relationship, secrecy.

Correspondence to: Mark McCann, MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, 200 Renfield Street, Glasgow G2 3QB, UK. E-mail: mark.mccann@glasgow.ac.uk

Submitted 1 May 2015; initial review completed 3 August 2015; final version accepted 30 November 2015

INTRODUCTION

More than 70% of European students have drunk alcohol at least once [1]. While alcohol has social functions, excess and early drinking may confer a risk of physical, emotional and social harm. While there may be no causal link between age of onset and later chronic alcohol problems [2,3], early use is predictive of future drinking patterns [4], which can contribute to long-term health outcomes [5–7], delinquency [8–10], mental health problems and risky sexual behaviour [11,12] and poor academic performance [8,13,14].

Providing emotional support for children is one element of the parental role [15]; another key element is parenting behaviours to minimize the risk of physical or social harm [16]. Parental monitoring—or parent's ongoing knowledge of their child's activities, whereabouts, etc.—has been proposed as the primary family-level determinant of alcohol outcomes: low parental monitoring has been associated with adolescent drinking outcomes [17–19]; one study showed that the importance of other family factors is negligible after accounting for parental knowledge [20]. Children choose whether or not to disclose information about their

activities and to keep secrets from their parents [21]. Frijns *et al.* reported that secrecy from parents was an important risk factor for adolescent psychosocial wellbeing and behavioural adjustment [22]; the child disclosure–adjustment link may actually reflect a secrecy–maladjustment link [23]. Adolescent agency to influence their environment (e.g. via selective information sharing with parents) may be particularly important in households where family relationships are already strained. Not only may parents change monitoring behaviour, but children may also choose different communication strategies depending on the quality of the relationship [24]. Few studies have been in a position to assess how alcohol relates to adolescent agency, or the extent to which young people influence their family environment actively.

While it is conceptually plausible that parent–child relationship could influence drinking (e.g. in response to stress [25] or hindered development of coping skills [15]), there is little evidence for a prospective association between parent–child relationship and drinking [26,27]. Relationship quality could influence monitoring patterns, and potentially the association between monitoring and child outcomes [28], so parental monitoring may be associated more strongly with alcohol use among children with poor family relationships and thus greater propensity to drink. By comparison, for adolescents with a more positive home environment, monitoring may not play as important a role.

Using longitudinal data across the period from early to mid-adolescence, this study will assess the relationships between parent–child attachment, parent and child behaviours influencing monitoring and adolescent alcohol use. It will also assess how associations differ according to home environment. Given the known gender differences in alcohol use [29] and in parent–child relationships [30,31], we will also assess differences by sex of the child. Our research questions are:

- 1 Does the quality of parent–child relationship and levels of parental monitoring affect alcohol use frequency?
- 2 Does alcohol use frequency influence subsequent levels of parental monitoring?
- 3 Does the monitoring and alcohol dynamic vary comparing high versus low attached families?
- 4 Do the associations change from early to late adolescence?

METHODS

This paper used data from the Belfast Youth Development Study, a longitudinal study of adolescent substance use. Between 2000 and 2011, children attending approximately 40 schools, colleges and special educational programmes across Northern Ireland were given questionnaires on a range of issues, including

substance use and family life. Pupils were in their first year of secondary school (approximately age 11) at the start of the study (academic year 2000/2001), and were surveyed annually until 2006/2007 (approximately age 17) whether or not in education, again approximately 10 years after the first sweep (2011). This report is based on data from the first 5 years of the study. There were 6156 pupils ever registered at participating schools during the study period. After excluding those who did not provide data at any time-point, withdrew from the study or those whose parents did not consent for them to participate, data were available for 5209 (85%) individuals. As this study focused on parent–child relationships, 195 children who stated that they did not live with their parents were removed from the analysis, leaving 5014 individuals. There were 77 individuals who did not complete the Inventory of Peer and Parental Attachment (IPPA) scale at any time-point. They were also dropped from the analysis, leaving 4937 individuals in the final analysis.

Study variables

Mental health was measured using the Strengths and Difficulties Questionnaire (SDQ) [32], a mental health screener for children and adolescents. The SDQ was completed in years 1 and 4. For family affluence, respondents were asked about the number of cars at their household, number of family holidays, parental employment, if they have their own bedroom, the type of house they lived in and eligibility for free school meals. Affluence measures were based on principal components analysis of these items [33]. Regarding living arrangements, respondents were asked with whom they lived, and responses were grouped into: both biological parents; step/foster-family (one biological and one step/foster-parent); and single-parent. Respondents in other households (predominantly siblings or grandparents) were excluded. Living arrangements in year 5 of the study were used in the analysis. If this information was missing in year 5, the previous year's data were used.

Analytical variables

Each year, participants were asked about how frequently they drank alcohol. Responses for each year were coded as: does not drink (0); rarely drinks (1); drinks monthly (2); and drinks weekly or more frequently [3]. Stattin & Kerr's (2000) measures of parental monitoring were asked in each year. Rather than looking at the overall monitoring measure, we focused on the mechanisms by which parents obtain information. The parental solicitation component enquired about how much parents ask their children about what they do; parental control relates to needing parents' permission or parental demands for information. While developed originally as a single scale, the child disclosure

component has been shown to comprise two factors [23]. The first centres on information offered to parents, referred to henceforth as 'disclosure', and the second centres on withholding information from parents, referred to as 'secrecy'. Higher scores on these scales indicated more 'positive' ratings, i.e. higher levels of parental control and solicitation, higher child disclosure or lower levels of secrecy (i.e. keeping fewer secrets). The inventory of peer and parental attachment [34] was also included in the analysis: this scale includes questions such as 'my parents respect my feelings' and 'I trust my parents'. The IPPA comprised 12 items on a three-point scale in year 1, while in years 3 and 4 it comprised 28 items on a five-point scale. Higher scores indicate better parent–child relationship. Respondents were split into high- and low-attached groups based on IPPA.

Statistical analysis

Cross-lagged path analysis assessed the association between alcohol use and monitoring dimensions over time. Monitoring and alcohol use were regressed on variables at a previous time-point. For alcohol use and monitoring this was always a 1-year lag; for SDQ and IPPA, the lag was to the last year they were asked (years 1, 3 or 4). The models contained the following components: (a) cross-sectional correlations between year 1 variables; (b) stability paths, the association between the same variable from one time-point to the next; and (c) cross-lagged paths, showing the association between one variable and subsequent levels of the other variables (see Fig. 1) These cross-lagged paths suggest to what extent alcohol use or monitoring is associated with subsequent change in the other construct, over and above that expected given their prior levels of drinking or monitoring. These models were fitted separately for each monitoring subcomponent; after this, the full model retaining all influential components was fitted. For this full model, subgroup analysis then estimated if stability and cross-lagged paths differed between (d) males and females and (e) high-attachment and low-attachment families. Finally, (f) the full cohort model was used to assess if the cross-lagged and stability paths differed across each study

year. All reported coefficients were conditioned on gender, affluence, SDQ, IPPA and living arrangements, and models were estimated using robust standard errors accounting for clustering at the school level; information was not collected on classes within schools. The Appendix contains further details on variables and the analyses of between-group differences. Latent profile analyses were conducted in MPlus 7, while the main analyses were performed using Stata version 13.1. The Sem and gsem routines were used in conjunction with mi (multiple imputation), while multiple group and time trend tests were conducted using mi testtransform; these results appear in the Supporting information Appendix. Results below are based on Stata's regress and ologit routines.

RESULTS

A total of 4937 were included into the study; 2348 (48%) were male and 2589 (52%) were female. Approximately 8–9% of the cohort showed some signs of mental health problems at both time-points that they were assessed. Approximately 71% (3545) of the cohort lived with both biological parents, 19% (942) lived with one parent only and approximately 9% (450) lived in a different household, e.g. parent and parent's partner, with foster-parents, etc. The mean [standard deviation (SD)] IPPA scores were 29.9 (3.8), 103.4 (21.8) and 102.9 (22.6) in years 1, 3 and 4, respectively.

A greater proportion of boys reported frequent drinking in year 1 (8% of males reported drinking weekly or more frequently compared to 3% of females), but in the following years there was little gender difference. Those living with step-parent or single-parent families drank more frequently from an earlier age than those with two biological parents, as did those with poorer mental health and less secure parental attachment.

Path models

Fit statistics that are appropriate to clustered data indicated reasonable model fit [35] (see Supporting information Appendix, Table A5). Tables 1–4 show the coefficients for




Figure 1 Representation of the path models resting the association between monitoring and alcohol use. Att = attachment, Alc = alcohol drinking frequency, Mon = parental monitoring

Table 1 Cross-lagged coefficients for the association between alcohol use frequency and parental control.

<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>
Year 2 Alcohol							
Year 1 Control	0.77 (0.72, 0.83)	Year 3 Alcohol		Year 4 Alcohol		Year 5 Alcohol	
Year Attachment	0.96 (0.89, 1.04)	Year 2 Control	0.76 (0.71, 0.81)	Year 3 Control	0.77 (0.72, 0.83)	Year 4 Control	0.84 (0.77, 0.92)
Year 2 Control		Year 1 Attachment	0.91 (0.85, 0.97)	Year 3 Attachment	0.965 (0.90, 1.04)	Year 4 Attachment	0.98 (0.89, 1.07)
Year 1 Alcohol		Year 3 Control		Year 4 Control		Year 5 Control	
None	Reference	Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol	
Rarely	-0.11 (-0.18, -0.04)	None	Reference	None	Reference	None	Reference
Monthly	-0.111 (-0.27, 0.04)	Rarely	-0.09 (-0.15, -0.03)	Rarely	-0.08 (-0.15, -0.01)	Rarely	-0.04 (-0.13, 0.06)
Weekly or more	-0.197 (-0.36, -0.03)	Monthly	-0.17 (-0.26, -0.08)	Monthly	-0.11 (-0.20, -0.03)	Monthly	-0.09 (-0.19, 0.01)
Year 1 Attachment	0.07 (0.04, 0.11)	Weekly or more	-0.262 (-0.35, -0.18)	Weekly or more	-0.18 (-0.26, -0.09)	Weekly or more	-0.19 (-0.30, -0.09)
		Year 1 Attachment	0.10 (0.06, 0.13)	Year 3 Attachment	0.05 (0.01, 0.08)	Year 4 Attachment	0.01 (-0.03, 0.04)

Coefficients for alcohol represent ordinal logistic regression odds ratios; coefficients for control represent linear regression standardized beta coefficients; coefficients adjusted for sex, living arrangements, affluence and mental health. CI = confidence interval. Coefficients in bold denote $P < 0.05$. Bold text denotes outcome variable.

Table 2 Cross-lagged coefficients for the association between alcohol use frequency and child disclosure.

<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>	<i>Path</i>	<i>Coefficient (95% CI)</i>
Year 2 Alcohol							
Year 1 Disclosure	0.77 (0.71, 0.83)	Year 3 Alcohol		Year 4 Alcohol		Year 5 Alcohol	
Year 1 Attachment	0.99 (0.92, 1.07)	Year 2 Disclosure	0.80 (0.74, 0.85)	Year 3 Disclosure	0.77 (0.71, 0.83)	Year 4 Disclosure	0.87 (0.80, 0.96)
Year 2 Disclosure		Year 1 Attachment	0.92 (0.86, 0.99)	Year 3 Attachment	1.04 (0.96, 1.12)	Year 4 Attachment	1.01 (0.91, 1.12)
Year 1 Alcohol		Year 3 Disclosure		Year 4 Disclosure		Year 5 Disclosure	
None	Reference	Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol	
Rarely	-0.15 (-0.22, -0.08)	None	Reference	None	Reference	None	Reference
Monthly	-0.29 (-0.45, -0.14)	Rarely	-0.10 (-0.16, -0.04)	Rarely	-0.13 (-0.22, -0.04)	Rarely	-0.12 (-0.21, -0.03)
Weekly or more	-0.22 (-0.39, -0.06)	Monthly	-0.17 (-0.25, -0.08)	Monthly	-0.24 (-0.34, -0.14)	Monthly	-0.23 (-0.33, -0.13)
Year 1 Attachment	0.13 (0.09, 0.17)	Weekly or more	-0.17 (-0.26, -0.08)	Weekly or more	-0.23 (-0.33, -0.14)	Weekly or more	-0.23 (-0.33, -0.13)
		Year 1 Attachment	0.13 (0.09, 0.16)	Year 3 Attachment	0.14 (0.11, 0.17)	Year 4 Attachment	0.19 (0.15, 0.23)

Coefficients for alcohol represent ordinal logistic regression odds ratios; coefficients for disclosure represent linear regression standardized beta coefficients; coefficients adjusted for sex, living arrangements, affluence and mental health. CI = confidence interval. Coefficients in bold denote $P < 0.05$. Bold text denotes outcome variable.

Table 3 Cross-lagged coefficients for the association between alcohol use frequency and child secrecy.

Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)
Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol		Year 5 Alcohol	
Year 1 Secrecy	0.71 (0.65, 0.76)	Year 2 Secrecy	0.77 (0.72, 0.83)	Year 3 Secrecy	0.73 (0.67, 0.79)	Year 4 Secrecy	0.81 (0.73, 0.90)
Year 1 Attachment	1.00 (0.92, 1.08)	Year 1 Attachment	0.91 (0.84, 0.97)	Year 3 Attachment	1.01 (0.93, 1.09)	Year 4 Attachment	1.02 (0.92, 1.12)
Year 2 Secrecy		Year 3 Secrecy		Year 4 Secrecy		Year 5 Secrecy	
Year 1 Alcohol		Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol	
None	Reference	None	Reference	None	Reference	None	Reference
Rarely	-0.21 (-0.29, -0.14)	Rarely	-0.21 (-0.28, -0.15)	Rarely	-0.19 (-0.26, -0.11)	Rarely	-0.33 (-0.42, -0.24)
Monthly	-0.32 (-0.48, -0.15)	Monthly	-0.33 (-0.43, -0.24)	Monthly	-0.31 (-0.40, -0.22)	Monthly	-0.47 (-0.57, -0.36)
Weekly or more	-0.41 (-0.57, -0.25)	Weekly or more	-0.40 (-0.50, -0.30)	Weekly or more	-0.39 (-0.48, -0.30)	Weekly or more	-0.54 (-0.65, -0.44)
Year 1 Attachment	0.09 (0.05, 0.13)	Year 1 Attachment	0.10 (0.06, 0.14)	Year 3 Attachment	0.14 (0.10, 0.17)	Year 4 Attachment	0.17 (0.13, 0.21)

Coefficients for alcohol represent ordinal logistic regression odds ratios; coefficients for secrecy represent linear regression standardized beta coefficients. Higher secrecy scores indicate lower levels of secret-keeping. Coefficients for disclosure represent linear regression standardized beta coefficients. Coefficients adjusted for sex, living arrangements, affluence and mental health. CI = confidence interval. Coefficients in bold denote $P < 0.05$. Bold text denotes outcome variable.

Table 4 Cross-lagged coefficients for the association between alcohol use frequency and parental solicitation.

Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)
Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol		Year 5 Alcohol	
Year 1 Solicitation	0.99 (0.92, 1.07)	Year 2 Solicitation	0.92 (0.86, 0.99)	Year 3 Solicitation	1.07 (0.99, 1.15)	Year 4 Solicitation	1.01 (0.93, 1.11)
Year 1 Attachment	0.91 (0.84, 0.98)	Year 1 Attachment	0.89 (0.83, 0.95)	Year 3 Attachment	0.89 (0.82, 0.96)	Year 4 Attachment	0.94 (0.85, 1.04)
Year 2 Solicitation		Year 3 Solicitation		Year 4 Solicitation		Year 5 Solicitation	
Year 1 Alcohol		Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol	
None	Reference	None	Reference	None	Reference	None	Reference
Rarely	-0.09 (-0.16, -0.02)	Rarely	0.01 (-0.06, 0.07)	Rarely	0.04 (-0.03, 0.11)	Rarely	0.05 (-0.05, 0.14)
Monthly	-0.20 (-0.36, -0.03)	Monthly	-0.00 (-0.10, 0.09)	Monthly	-0.00 (-0.10, 0.10)	Monthly	0.03 (-0.07, 0.13)
Weekly or more	-0.19 (-0.36, -0.02)	Weekly or more	-0.03 (-0.13, 0.06)	Weekly or more	0.01 (-0.08, 0.09)	Weekly or more	0.08 (-0.03, 0.18)
Year 1 Attachment	0.13 (0.09, 0.17)	Year 1 Attachment	0.13 (0.09, 0.17)	Year 3 Attachment	0.11 (0.08, 0.15)	Year 4 Attachment	0.07 (0.03, 0.10)

Coefficients for alcohol represent ordinal logistic regression odds ratios; coefficients for solicitation represent linear regression standardized beta coefficients; coefficients adjusted for sex, living arrangements, affluence and mental health; coefficients for disclosure represent linear regression standardized beta coefficients. CI = confidence interval. Coefficients in bold denote $P < 0.05$. Bold text denotes outcome variable.

Table 5 Cross-lagged coefficients for the association between alcohol use frequency, parental control, child disclosure and child secrecy.

Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)	Path	Coefficient (95% CI)
Year 2 Alcohol		Year 3 Alcohol		Year 4 Alcohol		Year 5 Alcohol	
Year 1 Control	0.83 (0.77, 0.90)	Year 2 Control	0.81 (0.75, 0.87)	Year 3 Control	0.83 (0.76, 0.90)	Year 4 Control	0.86 (0.78, 0.95)
Year 1 Disclosure	0.89 (0.82, 0.96)	Year 2 Disclosure	0.92 (0.85, 0.99)	Year 3 Disclosure	0.88 (0.80, 0.96)	Year 4 Disclosure	0.96 (0.86, 1.07)
Year 1 Secrecy	0.76 (0.70, 0.82)	Year 2 Secrecy	0.82 (0.76, 0.89)	Year 3 Secrecy	0.76 (0.70, 0.83)	Year 4 Secrecy	0.83 (0.75, 0.92)
Year 1 Attachment	1.06 (0.98, 1.15)	Year 1 Attachment	0.94 (0.88, 1.02)	Year 3 Attachment	1.10 (1.02, 1.19)	Year 4 Attachment	1.06 (0.95, 1.18)
Year 2 Control		Year 3 Control		Year 4 Control		Year 5 Control	
Year 1 Alcohol	Reference	Year 2 Alcohol	Reference	Year 3 Alcohol	Reference	Year 4 Alcohol	Reference
None	Reference	None	Reference	None	Reference	None	Reference
Rarely	-0.08 (-0.15, -0.01)	Rarely	-0.06 (-0.12, 0.01)	Rarely	-0.05 (-0.12, 0.02)	Rarely	-0.02 (-0.12, 0.07)
Monthly	-0.04 (-0.19, 0.12)	Monthly	-0.11 (-0.20, -0.02)	Monthly	-0.07 (-0.16, 0.03)	Monthly	-0.06 (-0.17, 0.04)
Weekly or more	-0.11 (-0.28, 0.06)	Weekly or more	-0.17 (-0.27, -0.08)	Weekly or more	-0.11 (-0.20, -0.02)	Weekly or more	-0.15 (-0.26, -0.04)
Year 1 Disclosure	0.06 (0.02, 0.10)	Year 2 Disclosure	0.06 (0.03, 0.10)	Year 3 Disclosure	0.08 (0.05, 0.13)	Year 4 Disclosure	0.07 (0.02, 0.11)
Year 1 Secrecy	0.06 (0.02, 0.10)	Year 2 Secrecy	0.05 (0.02, 0.09)	Year 3 Secrecy	0.05 (0.01, 0.08)	Year 4 Secrecy	0.03 (-0.01, 0.06)
Year 1 Attachment	0.04 (0.01, 0.08)	Year 1 Attachment	0.08 (0.04, 0.11)	Year 3 Attachment	-0.00 (-0.04, 0.03)	Year 4 Attachment	-0.03 (-0.07, 0.01)
Year 2 Disclosure		Year 3 Disclosure		Year 4 Disclosure		Year 5 Disclosure	
Year 1 Alcohol	Reference	Year 2 Alcohol	Reference	Year 3 Alcohol	Reference	Year 4 Alcohol	Reference
None	Reference	None	Reference	None	Reference	None	Reference
Rarely	-0.10 (-0.17, -0.03)	Rarely	-0.06 (-0.12, 0.01)	Rarely	-0.04 (-0.10, 0.03)	Rarely	-0.09 (-0.17, 0.00)
Monthly	-0.16 (-0.32, -0.00)	Monthly	-0.10 (-0.19, -0.01)	Monthly	-0.09 (-0.18, -0.00)	Monthly	-0.15 (-0.25, -0.06)
Weekly or more	-0.11 (-0.27, 0.06)	Weekly or more	-0.06 (-0.15, 0.04)	Weekly or more	-0.07 (-0.16, 0.02)	Weekly or more	-0.10 (-0.20, 0.00)
Year 1 Control	0.11 (0.07, 0.15)	Year 2 Control	0.12 (0.08, 0.15)	Year 3 Control	0.10 (0.06, 0.13)	Year 4 Control	0.02 (-0.02, 0.05)
Year 1 Secrecy	0.11 (0.07, 0.15)	Year 2 Secrecy	0.09 (0.05, 0.12)	Year 3 Secrecy	0.08 (0.05, 0.11)	Year 4 Secrecy	0.16 (0.12, 0.19)
Year 1 Attachment	0.10 (0.06, 0.14)	Year 1 Attachment	0.11 (0.08, 0.15)	Year 3 Attachment	0.10 (0.07, 0.14)	Year 4 Attachment	0.15 (0.11, 0.19)
Year 2 Secrecy		Year 3 Secrecy		Year 4 Secrecy		Year 5 Secrecy	
Year 1 Alcohol	Reference	Year 2 Alcohol	Reference	Year 3 Alcohol	Reference	Year 4 Alcohol	Reference
None	Reference	None	Reference	None	Reference	None	Reference
Rarely	-0.19 (-0.27, -0.12)	Rarely	-0.17 (-0.24, -0.11)	Rarely	-0.17 (-0.24, -0.09)	Rarely	-0.28 (-0.37, -0.19)
Monthly	-0.25 (-0.42, -0.08)	Monthly	-0.27 (-0.37, -0.17)	Monthly	-0.27 (-0.37, -0.18)	Monthly	-0.38 (-0.48, -0.28)
Weekly or more	-0.35 (-0.51, -0.18)	Weekly or more	-0.30 (-0.41, -0.20)	Weekly or more	-0.33 (-0.43, -0.24)	Weekly or more	-0.42 (-0.53, -0.32)
Year 1 Control	0.06 (0.02, 0.10)	Year 2 Control	0.04 (0.00, 0.07)	Year 3 Control	0.02 (-0.01, 0.06)	Year 4 Control	0.06 (0.02, 0.09)
Year 1 Disclosure	0.07 (0.03, 0.11)	Year 2 Disclosure	0.16 (0.12, 0.20)	Year 3 Disclosure	0.11 (0.07, 0.15)	Year 4 Disclosure	0.24 (0.19, 0.28)
Year 1 Attachment	0.07 (0.03, 0.11)	Year 1 Attachment	0.06 (0.02, 0.10)	Year 3 Attachment	0.09 (0.05, 0.13)	Year 4 Attachment	0.07 (0.03, 0.11)

Coefficients for alcohol represent ordinal logistic regression odds ratios; coefficients for control, disclosure and secrecy represent linear regression standardized beta coefficients; coefficients for sex, living arrangements, affluence and mental health, CI = confidence interval. Coefficients in bold denote $P < 0.05$. Bold text denotes outcome variable.

the bidirectional associations between alcohol use and parental control, child disclosure, child secrecy and parental solicitation, respectively. Parental disclosure showed little association with alcohol use and was dropped from further models. Table 5 shows the mutually adjusted model testing cross-lagged paths between alcohol, parental control, child disclosure and secrecy. Tables showing tests for time trends, gender differences and parental attachment group differences are reported in the Supporting information Appendix.

Influences on frequency of alcohol use

Apart from the parental solicitation subscale, higher levels of parental monitoring were associated with less frequent use of alcohol. When observing a 1 SD increase in parental monitoring in year 4, there was a lower odds of frequent drinking in year 5; this was observed for parental control, beta 0.84 [95% confidence interval (CI) = 0.77, 0.92], child disclosure, beta 0.87 (95% CI = 0.80, 0.96) and child secrecy, beta 0.81 (95% CI = 0.73, 0.90), but there was no change in drinking frequency associated with levels of parental solicitation (beta 1.01, 95% CI = 0.93, 1.11). There was no evidence of variation over time for the influence of parental control ($P = 0.29$) and child disclosure ($P = 0.53$) on alcohol use. The association between child secrecy and alcohol use did appear to vary over time ($P = 0.02$), with less of an association with the secrecy scale in year 1 ($P = 0.03$) than the scale in year 2 ($P = 0.47$). IPPA showed little association with drinking frequency [odds ratio (OR) = 0.94, 95% CI = 0.85, 1.04] and the influence of monitoring did not vary comparing high and low IPPA groups.

Influences on parental control

More frequent adolescent alcohol use was associated with lower parental control. Respondents drinking weekly or more frequently showed an approximately 0.20 SD reduction in parental control in the subsequent year. This association did not appear to vary over time ($P = 0.18$). Respondents with higher disclosure scores reported slightly higher subsequent parental control, and this did not appear to vary over time ($P = 0.96$); a 1-unit increase in year 4 disclosure was associated with a 7% increase in parental control (beta 0.07, 95% CI = 0.02, 0.11). Lower levels of child secrecy were associated with higher parental control in the earlier years of the study, but by age 14 the association attenuated to suggest no effect. There was some evidence for a time trend ($P = 0.09$). Parental attachment did not appear to influence parental control (year 4 IPPA standardized beta 0.028, 95% CI = -0.07, 0.01).

Influences on child disclosure

Alcohol use was associated with lower levels of child disclosure. Compared to non-drinkers, year 5 disclosure scores were lower for infrequent (beta -0.09, 95% CI = -0.17, 0.00), monthly (beta -0.15, 95% CI = -0.25, -0.06) and weekly (beta -0.10, 95% CI = -0.20, 0.00) drinkers. There was little association of child disclosure with parental control (beta 0.02, 95% CI = 0.02, 0.05), but more of an association with child secrecy (beta 0.16, 95% CI = 0.12, 0.19) and parental attachment (beta 0.15, 95% CI = 0.11, 0.19). The association with alcohol ($P = 0.17$) did not vary over time, while the control-disclosure association did ($P < 0.001$), with stronger evidence of a change in association in the later years of the study ($P = 0.09$), possibly explained by there being little association in year 5. The secrecy-disclosure association also varied over time ($P < 0.001$), again suggesting greater change at the later years.

Influences on child secrecy

The strongest association found across any of the models was the association between alcohol use and subsequent levels of child secrecy. Compared to non-drinkers in year 4, the rates of secrecy were higher in year 5: for infrequent drinkers by beta 0.28 (95% CI = 0.37, 0.19), monthly drinkers, beta 0.38 (95% CI = 0.48, 0.28) and weekly drinkers, beta 0.42 (95% CI = 0.53, 0.32). Alcohol use was, in fact, a stronger predictor of subsequent secrecy-keeping than previous levels of secrecy (beta 0.27, 95% CI = 0.23, 0.31). Higher parental control (beta 0.06, 95% CI = 0.02, 0.09), parent-child attachment (beta 0.07, 95% CI = 0.03, 0.11) and child disclosure (beta 0.24, 95% CI = 0.19, 0.28) were also associated with lower levels of secrecy. The association between disclosure and secrecy appeared to vary over time ($P < 0.001$), with evidence for a change in association between each time-point. There was no evidence of time trends for alcohol ($P = 0.33$), control ($P = 0.14$) or attachment ($P = 0.29$).

Variation by gender

Based on tests that the difference in the coefficients between the male and female models was equal to zero, the strongest suggestion of a gender difference was found for the alcohol-parental control association ($P = 0.08$), indicating that alcohol use was associated with a greater reduction in parental control among boys than girls.

Variation by high versus low parental attachment

There was no evidence of variation between high and low IPPA groups, except for differences in how alcohol use relates to subsequent parental control ($P = 0.04$) and child

disclosure ($P = 0.03$). The difference in parental control appeared greatest at older ages (year 5 $P = 0.01$), but at the youngest ages for child disclosure ($P = 0.03$). Alcohol use is followed by a greater reduction in parental control, but a lesser reduction in disclosure among the securely compared to the less securely attached group.

DISCUSSION

The study demonstrated the association between parental monitoring and adolescent alcohol use. The findings correspond with previous research [17,36], and additionally highlight the specific roles of parental control, children's withholding information and a reverse causal influence of drinking to reduce levels of monitoring. Parental monitoring and child drinking appear to influence each other. This suggests that interventions targeting changes in parenting practice and child drinking as dynamic processes could have benefits over and above preventing or delaying alcohol use initiation. There was little evidence of differences in these associations by gender or parent-child relationship, and the influence of monitoring on alcohol use was consistent over time.

Alcohol use seemed to have a stronger influence on parental control than on child disclosure, despite the fact that adolescents directly influence their own behaviour, but not that of their parents. There are a number of explanations: (1) children who drink more often may feel less controlled by their parents; (2) age may confound the control-alcohol use relationship; (3) drinking adolescents may have greater incentive to negotiate autonomy and independence, to facilitate socializing in drinking environments; and (4) parents are aware of their children's alcohol use and reduce control as a result. Point 1 does not account for the fact that the control-alcohol use relationship remains after accounting for levels of secrecy. Point 2 can be ruled out, because the association between drinking and control occurs at all ages. Points 3 and 4 attest to early identification of drinking. Alcohol initiation may lead to a child-directed process; alternatively, knowledge of a child's drinking may indicate independence and maturity or highlight the futility of the parent's efforts to control behaviour [37]. While parents may wish to facilitate adolescent self-determination, unsupported adolescent decision-making could actually worsen wellbeing [38]; this issue is particularly salient, given social pressures to drink [39,40], and given that this relationship is present from age 11. Teenagers in 2015 are entering a recommodified alcohol market [41], comparable in social environment to that of their parents, but markedly different in drinking intensity. Parents may endorse drinking autonomy at the same age as was afforded to them, but in so doing are facilitating entry into a markedly different risk environment. Differentially supporting autonomy for identity development in

some social arenas, while maintaining control and parental authority in relation to risk issues such as drinking, may represent 'optimal' support for adolescent development [42].

Parental solicitation showed little association with alcohol outcomes [43]. It may be easier to encourage parental behaviour change in terms of initiating conversation than it is to change behaviour surrounding curfews or setting boundaries on adolescent behaviour and whereabouts [44]. Parental attachment had little direct influence on alcohol use or parental behaviour; any cross-sectional associations are not borne out in longitudinal data [28], suggesting that this is a non-causal association. However, attachment interacts with the reverse causal process whereby alcohol use affects the home environment. A recent model from family systems theory outlined how sibling and parent relationships can lead to substance use problems [45]. Understanding a reverse causal process, and how this process differs according to characteristics of the family system, could have implications for integrating external social influences into family therapeutic work.

Considering the developmental and social differences with age, there was surprisingly little evidence for differences in younger or older adolescence. There is a general shift in the importance of family and peers from childhood to adolescence [46], but our results suggest that the role of parents in determining alcohol behaviour is consistently important. A study of parents uncovered concerns about the school and peer context—rather than the family home—as the best place to learn responsible drinking [47]; this suggests that parents are receptive to implementing a preventive approach to health risk behaviours. As such, in addition to school, family is an appropriate context in which to deliver preventive interventions [48,49].

A major strength of the study is its large sample size, providing sufficient power to test for group differences and changes over time. None the less, this study has several limitations. Data were collected from children, so we do not know if parents knew, suspected or permitted their children to drink. Furthermore, school environment may influence alcohol use, and this study provides no insight into the interplay between home and school contexts. Future research could assess in greater detail how characteristics of school interact with family processes to influence drinking behaviour.

Variation in parenting practices may amplify inequalities in life-style-related illnesses or social harms in later life, particularly given the socio-economic patterning in parenting approaches [50]. The family environment is thus a potential arena for preventive interventions and policy, but not simply from the traditional social service model of intervening with 'at-risk' families [51]. The reverse causal processes uncovered in this study highlight the fact that

parenting behaviour relates to dynamic family processes, rather than fixed parenting skills and knowledge. Preventive interventions which conceptualize parenting as a modifiable, resource-based practice [52] may have benefits for family functioning and alcohol outcomes. Given the prevalence of alcohol use in adolescence and among the wider population, broader, resource-based approaches may realize greater benefits than targeted interventions [53].

Research often endorses the early years as the key target for successful health [54] and social wellbeing [55] policies, advice that has been picked up at the state level, certainly in the United Kingdom [56]; family interventions for adolescents have received somewhat less attention. Given that adolescence is the critical period for the beginning of alcohol use, and that the harms of excessive drinking are not confined to children from 'problem' families, support for adolescent parenting—rather than alcohol awareness for parents—may be a fruitful target for public policy relating to young people's health risk behaviour.

Declaration of interests

None.

Acknowledgements

Many thanks to all of the schools and individuals for taking part in the Belfast Youth Development Study. The Belfast Youth Development Study was funded by the Research and Development Division of the Northern Ireland Health and Social Care Public Health Agency. This research was funded by a grant from Alcohol Research UK (Grant reference: R 2011/06). M.McC. holds a Medical Research Council/University fellowship supported by MRC partnership grant MC/PC/13 027.

References

- Hibell B., Guttormsson U., Ahlström S., Balakireva O., Bjarnason T., Kokkevi A. *et al.* The 2011 ESPAD report. Substance use among students among students in 36 European countries. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN); 2012.
- Maimaris W., McCambridge J. Age of first drinking and adult alcohol problems: systematic review of prospective cohort studies. *J Epidemiol Community Health* 2014; **68**: 268–74.
- Prescott C. A., Kendler K. S. Age at first drink and risk for alcoholism: a noncausal association. *Alcohol Clin Exp Res* 1999; **23**: 101–7.
- Irons D. E., Iacono W. G., McGue M. Tests of the effects of adolescent early alcohol exposures on adult outcomes. *Addiction* 2015; **110**: 269–78.
- Probst C., Roerecke M., Behrendt S., Rehm J. Socioeconomic differences in alcohol-attributable mortality compared with all-cause mortality: a systematic review and meta-analysis. *Int J Epidemiol* 2014; **43**: 1314–27.
- Patra J., Taylor B., Irving H., Roerecke M., Baliunas D., Mohapatra S. *et al.* Alcohol consumption and the risk of morbidity and mortality for different stroke types—a systematic review and meta-analysis. *BMC Public Health* 2010; **10**: 258–2458-10-258.
- Roerecke M., Rehm J. Irregular heavy drinking occasions and risk of ischemic heart disease: a systematic review and meta-analysis. *Am J Epidemiol* 2010; **171**: 633–44.
- Peleg-Oren N., Saint-Jean G., Cardenas G. A., Tammara H., Pierre C. Drinking alcohol before age 13 and negative outcomes in late adolescence. *Alcohol Clin Exp Res* 2009; **33**: 1966–72.
- Felson R. B., Teasdale B., Burchfield K. B. The influence of being under the influence alcohol effects on adolescent violence. *J Res Crime Delinquency* 2008; **45**: 119–41.
- Ellickson P. L., Tucker J. S., Klein D. J. Ten-year prospective study of public health problems associated with early drinking. *Pediatrics* 2003; **111**: 949–55.
- Fergusson D. M., Lynskey M. T., Horwood L. J. Factors associated with continuity and changes in disruptive behavior patterns between childhood and adolescence. *J Abnorm Child Psychol* 1996; **24**: 533–53.
- Cavazos-Rehg P. A., Spitznagel E. L., Bucholz K. K., Nurnberger J. Jr., Edenberg H. J., Kramer J. R. *et al.* Predictors of sexual debut at age 16 or younger. *Arch Sex Behav* 2010; **39**: 664–73.
- Balsa A. I., Giuliano L. M., French M. T. The effects of alcohol use on academic achievement in high school. *Econ Educ Rev* 2011; **30**: 1–15.
- Barry A. E., Chaney B., Chaney J. D. The impact of truant and alcohol-related behavior on educational aspirations: a study of US high school seniors. *J School Health* 2011; **81**: 485.
- Bowlby J. *A Secure Base: Parent–Child Attachment and Healthy Human Development*. New York: Basic Books; 2008.
- Asmussen K., Corlyon J., Hauari H., La Placa V. Supporting Parents of Teenagers. 2007; RR830.
- Fosco G. M., Stormshak E. A., Dishion T. J., Winter C. E., Fosco G. M., Stormshak E. A. *et al.* Family Relationships and Parental Monitoring During Middle School as Predictors of Early Adolescent Problem Behavior. *J Clin Child Adolesc Psychol* 2012; **41**: 202.
- Bremner P., Burnett J., Nunney E., Ravat M., Mistral W. *Young People, Alcohol and Influences: A Study of Young People and Their Relationship With Alcohol*. York: Joseph Rowntree Foundation; 2011.
- Ryan S. M., Jorm A. F., Lubman D. I. Parenting factors associated with reduced adolescent alcohol use: a systematic review of longitudinal studies. *Aust NZ J Psychiatry* 2010; **44**: 774–83.
- Ledoux S., Miller P., Choquet M., Plant M. Human study. Family structure, parent–child relationships, and alcohol and other drug use among teenagers in France and the United Kingdom. *Alcohol Alcohol* 2002; **37**: 52–2.
- Keijsers L., Laird R. D. Introduction to special issue. Careful conversations: adolescents managing their parents' access to information. *J Adolesc* 2010; **33**: 255–9.
- Frijns T., Finkenauer C., Vermulst A. A., Engels R. C. Keeping secrets from parents: longitudinal associations of secrecy in adolescence. *J Youth Adolesc* 2005; **34**: 137–48.
- Frijns T., Keijsers L., Branje S., Meeus W. What parents don't know and how it may affect their children: qualifying the disclosure–adjustment link. *J Adolesc* 2010; **33**: 261–70.
- Mazur M. A., Ebesu Hubbard A. S. 'Is there something I should know?': topic avoidant responses in parent–adolescent communication. *Communication Rep* 2004; **17**: 27–37.

25. Repetti R. L., Taylor S. E., Seeman T. E. Risky families: family social environments and the mental and physical health of offspring. *Psychol Bull* 2002; **128**: 330.
26. Visser L., de Winter A. F., Reijneveld S. A. The parent–child relationship and adolescent alcohol use: a systematic review of longitudinal studies. *BMC Public Health* 2012; **12**: 886.
27. Chan G. C., Kelly A. B., Toumbourou J. W., Hemphill S. A., Young R. M., Haynes M. A. *et al.* Predicting steep escalations in alcohol use over the teenage years: age-related variations in key social influences. *Addiction* 2013; **108**: 1924–32.
28. van der Vorst H., Engels R. C. M. E., Meeus W., Dekovic M., Vermulst A. Parental attachment, parental control, and early development of alcohol use: a longitudinal study. *Psychol Addict Behav* 2006; **20**: 107–16.
29. Schulte M. T., Ramo D., Brown S. A. Gender differences in factors influencing alcohol use and drinking progression among adolescents. *Clin Psychol Rev* 2009; **29**: 535–47.
30. Eccles J. S., Jacobs J. E., Harold R. D. Gender role stereotypes, expectancy effects, and parents' socialization of gender differences. *J Soc Issues* 1990; **46**: 183–201.
31. Starrels M. E. Gender differences in parent–child relations. *J Fam Issues* 1994; **15**: 148–65.
32. Goodman A., Goodman R. Strengths and difficulties questionnaire as a dimensional measure of child mental health. *J Am Acad Child Adolesc Psychiatry* 2009; **48**: 400–3.
33. Vyas S., Kumaranayake L. Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plan* 2006; **21**: 459–68.
34. Armsden G. C., Greenberg M. T. The inventory of parent and peer attachment: Individual differences and their relationship to psychological well-being in adolescence. *J Youth Adolesc* 1987; **16**: 427–54.
35. Hooper D., Coughlan J., Mullen M. Structural equation modelling: guidelines for determining model fit. *Articles* 2008; **2**.
36. Bergh D., Hagquist C., Starrin B. Parental monitoring, peer activities and alcohol use: A study based on data on Swedish adolescents. *Drugs Educ Prev Policy* 2011; **18**: 100–7.
37. Dishion T. J., Nelson S. E., Bullock B. M. Premature adolescent autonomy: parent disengagement and deviant peer process in the amplification of problem behaviour. *J Adolesc* 2004; **27**: 515–30.
38. Van Petegem S., Beyers W., Vansteenkiste M., Soenens B. On the association between adolescent autonomy and psychosocial functioning: examining decisional independence from a self-determination theory perspective. *Dev Psychol* 2012; **48**: 76.
39. Winpenny E. M., Marteau T. M., Nolte E. Exposure of children and adolescents to alcohol marketing on social media websites. *Alcohol Alcohol* 2014; **49**: 154–9.
40. Leung R. K., Toumbourou J. W., Hemphill S. A. The effect of peer influence and selection processes on adolescent alcohol use: a systematic review of longitudinal studies. *Health Psychol Rev* 2014; **8**: 426–57.
41. Measham E., Brain K. 'Binge' drinking, British alcohol policy and the new culture of intoxication. *Crime Media Cult* 2005; **1**: 262–83.
42. Koepke S., Denissen J. J. Dynamics of identity development and separation–individuation in parent–child relationships during adolescence and emerging adulthood—a conceptual integration. *Dev Rev* 2012; **32**: 67–88.
43. Keijsers L. Developmental links of adolescent disclosure, parental solicitation, and control with delinquency: moderation by parental support. *Dev Psychol* 2009; **45**: 1314.
44. Higgins K., Macdonald G., McLaughlin K., O'Hara L., McCann M., Moriarty J. *Parenting UR Teen. A Randomised Trial of Implementation and Effectiveness. End of Study Report (June 2012)*. Belfast: Institute of Child Care Research, Queen's University Belfast; 2012.
45. Feinberg M. E., Solmeyer A. R., McHale S. M. The third rail of family systems: sibling relationships, mental and behavioral health, and preventive intervention in childhood and adolescence. *Clin Child Fam Psychol Rev* 2012; **15**: 43–57.
46. Nickerson A. B., Nagle R. J. Parent and peer attachment in late childhood and early adolescence. *J Early Adolesc* 2005; **25**: 223–49.
47. Valentine G., Gould M., Keenan J. *Alcohol Consumption and Family Life*. York: Joseph Rowntree Foundation; 2010. Web. 13 Jan. 2016.
48. Gates S., McCambridge J., Smith L. A., Foxcroft D. Interventions for prevention of drug use by young people delivered in non-school settings. *Cochrane Database Syst Rev* 2006;(1): CD005030.
49. Foxcroft D. R., Tsertsvadze A. Cochrane review: universal school-based prevention programs for alcohol misuse in young people. *Evid Based Child Health* 2012; **7**: 450–575.
50. Odgers C. L., Caspi A., Russell M. A., Sampson R. J., Arseneault L., Moffitt T. E. Supportive parenting mediates neighborhood socioeconomic disparities in children's antisocial behavior from ages 5 to 12. *Dev Psychopathol* 2012; **24**: 705–21.
51. Dingwall R., Eekelaar J., Murray T. *The Protection of Children: State Intervention and Family Life*. New Orleans, LA: Quid Pro Books; 2014.
52. Daly M. Parenting support policies in Europe. *Fam Relat Soc* 2013; **2**: 159–74.
53. Rose G. *The Strategy of Preventive Medicine*. Oxford: Oxford University Press; 1992.
54. Oberklaid E., Baird G., Blair M., Melhuish E., Hall D. Children's health and development: approaches to early identification and intervention. *Arch Dis Child* 2013; **98**: 1008–11.
55. Waldfogel J., Washbrook E. Early years policy. *Child Dev Res* 2011; **2011**: 343016.
56. Allen G. *Early Intervention: The Next Steps. An independent report to Her Majesty's Government*. London: TSO for the Cabinet Office; 2011.

Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site:

Table A1 Number of respondents and cohort characteristics across five years

Table A2 Time-varying characteristics across five years

Table A3 Frequency of alcohol use across five years by gender

Table A4 Mean (sd.) for parental monitoring subscales by year and gender

Table A5 Fit statistics for Structural Equation Models using linear alcohol measures

Table A6 Cross lagged coefficients for the association between alcohol use frequency and parental control: linear alcohol use measure

Table A7 Cross lagged coefficients for the association between alcohol use frequency and child disclosure: linear alcohol use measure

Table A8 Cross lagged coefficients for the association between alcohol use frequency and child secrecy: linear alcohol use measure

Table A9 Cross lagged coefficients for the association between alcohol use frequency and parental solicitation: linear alcohol use measure

Table A10 Cross lagged coefficients for the association between alcohol use frequency, parental control, child

disclosure and child secrecy: linear alcohol use measure

Table A11 Tests of difference in estimated coefficients over time

Table A12 Difference between male and female model coefficients, tested jointly and within single years

Table A13 Difference between coefficients comparing high attachment and low attachment groups, tested jointly and within single years